

Robert P. Bruce, MD

*Corporate Medical Plaza,
Building 1
10701 Nall Ave., Suite 200
Overland Park, KS 66211*


*Lee's Summit Office
3151 NE Carnegie Dr.,
Suite B
Lee's Summit, MO 6406*

*Olathe Office
20920 W 151st St., Suite 100
Olathe, KS 66061*

Office (913)381-5225

If you have any questions or concerns, please call.

Rotator Cuff Repair



Pre-Operative Instructions

- ✓ **Do not eat or drink after midnight the night before surgery.** This includes chewing gum, candy and mints. No smoking or use of tobacco products.
- ✓ Stop all blood thinners 1 week prior to surgery, unless directed otherwise. This includes coumadin, aspirin, and anti-inflammatories such as ibuprofen, Nuprin, or Aleve. You may use acetaminophen for pain.
- ✓ Stop all herbal supplements 1 week prior to surgery.
- ✓ Plan to spend approximately 4 hours at the surgical facility.
- ✓ Arrange for someone to drive you home and stay with you through the night and for the first 24 hours after surgery.
- ✓ Set up your post-op appointment for 7-10 days after surgery at 913-381-5225 if you have not already done so. (You may also request an appointment on our patient portal)

Robert P. Bruce, MD

Post-Operative Instructions

- ✓ Apply ice to the shoulder for 20 minutes each hour for the first 2 days, then as frequently as needed to decrease swelling and for comfort. You should use ice after therapy.
- ✓ Keep the dressing dry and in place for 3 days. On the 3rd day you may apply a new dry gauze dressing. **Do not** get the incision wet.
- ✓ You will have a shoulder immobilizer in place after surgery. You may remove your hand and arm from the immobilizer only to perform the exercises described in this brochure and for daily hygiene.
- ✓ All motion for your affected arm during the immediate post-operative period will be **passive**, meaning that the movement is not produced by your shoulder muscles. Instead the forces of gravity or momentum will provide the movement. Do not attempt to raise your arm away from your body at this time.
- ✓ Perform the range of motion exercises 2-3 times per day. Begin these the day of surgery. When you come in for your post-op visit Dr. Bruce may order formal physical therapy (PT).
- ✓ You may do range of motion at the elbow and wrist as tolerated, but do not hold any weight when doing so.
- ✓ Swelling in your affected hand and forearm is not uncommon. You may elevate the lower arm and squeeze a small ball frequently to relieve this. You may also notice bruising in the arm below the incision.
- ✓ Take the anti-inflammatory medication as prescribed to you.
- ✓ If you have not received a regional block, take the pain medication regularly for the first 48 hours, then as needed.
- ✓ If you received a regional block to help with post-op pain control and have not taken any oral pain medication throughout the day, go ahead and take the pain medication before you go to bed.
- ✓ Be aware that a common side effect of the pain medications prescribed to you is constipation. You should remain well hydrated and include fiber in your diet. If necessary, you may use an over the counter stool softener or laxative.
- ✓ Start slowly with your diet, progressing from clear liquids to a regular diet. Avoid heavy or greasy foods the first 24 hours after surgery.
- ✓ You may find you are more comfortable sleeping in a recliner for the first few days.

PASSIVE PENDULUM EXERCISES

Bend forward 90 degrees at the waist, using a table for support. Rock your body in a circular pattern to move the arm clockwise 10 times, then counterclockwise 10 times. Let gravity and momentum provide the movement.

DO NOT USE YOUR SHOULDER MUSCLES TO MOVE YOUR ARM!

